

# Target Corporation

**-PE Product Labels –**

**-continued**

NDC 11673-410-06

Compare to active ingredients in **Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold\***

**maximum strength  
daytime  
severe cold**

**acetaminophen**  
(pain reliever/fever reducer)  
**dextromethorphan HBr** (cough suppressant)  
**guaifenesin** (expectorant)  
**phenylephrine HCl** (nasal decongestant)

relieves aches, fever and sore throat  
controls cough  
relieves nasal and chest congestion  
thins and loosens mucus

**up&up**

AGES  
**12+**  
YEARS

6 FL OZ (177 mL) PLD-83468 LB004227

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY  
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

094 14 0185 R01  
C-000274-01-049  
Dist. by Target Corp.  
Minneapolis, MN 55403  
TM & ©2021 Target Brands, Inc.

PLD-83468  
LB004228

PEEL CORNER FOR DRUG FACTS ▲

## Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

## Uses

- temporarily relieves these common cold and flu symptoms
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe ►

## Drug Facts (continued)

skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

PEEL CORNER FOR MORE DRUG FACTS ▲

## Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ►

## Drug Facts (continued)

## Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

## Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

## Questions or comments?

Call 1-800-910-6874

## Children's Mucus Multi-Symptom Cold Relief

**Drug Facts**

**Active ingredients (in each 5 mL)**

Dextromethorphan HBr 5 mg	Cough suppressant
Guafenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

**Uses** ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help your child get to sleep ■ nasal congestion due to a cold ■ stuffy nose

**Warnings**  
Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if your child has** ■ heart disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with asthma

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur ■ symptoms do not get better within 7 days or occur with fever ■ cough lasts more than 7 days, comes back, or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ do not take more than 6 doses in any 24-hour period ■ use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device. ■ keep dosing cup with product ■ dose as follows or as directed by a doctor ■ mL = milliliter

Age	Dose
Children 6 years to under 12 years	10 mL every 4 hours
Children 4 years to under 6 years	5 mL every 4 hours
Children under 4 years	do not use

**Other information** ■ each 5 mL contains sodium 5 mg ■ store between 20-25°C (68-77°F). Do not refrigerate

## Children's Mucus Multi-Symptom Cold Relief (continued)

**Drug Facts (continued)**

**Inactive ingredients** citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-800-910-8874

**Children's Nighttime Multi-Symptom Cold Relief**

**Drug Facts**

**Active ingredients (in each 10 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** ■ temporarily relieves these common cold and flu symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ sneezing ■ runny nose ■ nasal congestion ■ cough ■ combats cough to help your child get to sleep ■ temporarily reduces fever

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

In a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. ■ with any other drug containing diphenhydramine even when used on the skin ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product ■ to make a child sleepy

## Children's Nighttime Multi-Symptom Cold Relief (cont.)

**Drug Facts (continued)**

**Ask a doctor before use if your child has** ■ liver disease ■ heart disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ glaucoma ■ a breathing problem such as chronic bronchitis ■ persistent or chronic cough such as occurs with asthma ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if your child is** ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

**When using this product** ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days ■ indigestion or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not give more than 5 doses in any 24-hour period ■ if needed, repeat dose every 4 hours until symptoms last ■ do not give more than 5 days unless directed by a doctor ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter ■ dose as follows or as directed by a doctor ■ Children 6 to under 12 years of age: 10 mL every 4 hours ■ Children under 6 years of age: do not use

**Other information** ■ each 10 mL contains sodium 6 mg ■ store between 20-25°C (68-77°F). Do not refrigerate

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-800-910-8874

## Compare to active ingredients in Children's Mucinex® Multi-Symptom Cold\*

## children's day time multi-symptom cold relief

dextromethorphan HBr 5 mg  
(cough suppressant)  
guaifenesin 100 mg  
(expectorant)  
phenylephrine HCl 2.5 mg  
(nasal decongestant)

relieves stuffy nose  
controls cough  
chest congestion  
breaks up mucus  
alcohol free



4 FL OZ (118 mL)  
8 FL OZ (236 mL) TOTAL

## Compare to active ingredients in Children's Mucinex® Night Time Multi-Symptom Cold\*

## children's night time multi-symptom cold relief

acetaminophen 325 mg  
(pain reliever/fever reducer)  
diphenhydramine HCl 12.5 mg  
(antihistamine/cough suppressant)  
phenylephrine HCl 5 mg  
(nasal decongestant)

relieves stuffy nose  
controls cough  
runny nose & sneezing  
fever & sore throat  
alcohol free



4 FL OZ (118 mL)

NDC 11673-420-08

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold and Children's Mucinex® Multi-Symptom Cold.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org



100% satisfaction guaranteed or your money back.

094 14 0025 R02  
100-000021-01-193  
Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2019 Target Brands, Inc.



7 15256 33607 9

PLD-CS43A  
PLD-CS44A  
FC005821

Lot No.:  
Exp. Date:



NDC 11673-460-06

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu\*

# maximum strength nighttime cold and flu

**acetaminophen**

(pain reliever/fever reducer)

diphenhydramine HCl

(antihistamine/cough suppressant)

phenylephrine HCl (nasal decongestant)

relieves aches, fever and sore throat

controls cough

relieves nasal congestion

relieves runny nose and sneezing



AGES  
**12+**  
YEARS

6 FL OZ (177 mL)

PLD-B377D LB004229

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu.

094 14 0185

C-000274-01-049

Dist. by Target Corp.

Mpls., MN 55403

TM & ©2022 Target Brands, Inc.

PLD-B377D

LB004230

## Drug Facts

### Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Diphenhydramine HCl 25 mg.....Antihistamine/cough suppressant  
Phenylephrine HCl 10 mg.....Nasal decongestant

### Purposes

PEEL CORNER FOR MORE DRUG FACTS

## Drug Facts (continued)

### Uses

- temporarily relieves these common cold and flu symptoms
- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- runny nose
- sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen

## Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

PEEL CORNER FOR MORE DRUG FACTS

## Drug Facts (continued)

### When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Drug Facts (continued)

### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

### Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

### Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Questions or comments?

Call 1-800-910-6874

**congestion, non-drowsy**  
**sinus PE**

**Drug Facts** (continued)  
If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.  
■ children under 12 years: ask a doctor

**Other information**  
■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**  
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ see end flap for expiration date and lot number

**Inactive ingredients** croscarmellose sodium, dextrose monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40, lecithin, magnesium stearate, malto-dextrin, microcrystalline cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium citrate dihydrate, titanium dioxide

**Questions?** Call 1-800-910-6874

**congestion, non-drowsy**  
**sinus PE**

**Drug Facts**  
**Active Ingredient (in each tablet)** Purpose  
Phenylephrine HCl 10 mg ..... Nasal decongestant

**Uses**  
■ temporarily relieves sinus congestion and pressure  
■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

**Warnings**  
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
■ heart disease ■ diabetes  
■ thyroid disease ■ high blood pressure  
■ difficulty in urination due to enlargement of the prostate gland

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor if**  
■ symptoms do not improve within 7 days or occur with fever  
■ nervousness, dizziness, or sleeplessness occur

**congestion, non-drowsy**  
**sinus PE**

**36 TABLETS (10 mg EACH)**

**up&up**

**36 TABLETS**

**3 11673 45307 0**

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**congestion, non-drowsy**  
**sinus PE**

**Compare to active ingredient in Sudafed PE® Sinus Congestion\***

**congestion, non-drowsy**  
**sinus PE**

phenylephrine HCl tablets, nasal decongestant

sinus pressure and congestion  
maximum strength  
pseudoephedrine free

**up&up**

**36 TABLETS (10 mg EACH)**

**up&up**

**36 TABLETS**

**3 11673 45307 0**

NDC 11673-453-07

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

094 03 0434 R00 C-001472-01-109  
Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2021 Target Brands, Inc.

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE® Sinus Congestion.  
50844 ORG082045307

B-1920-453-07  
ORG082045307

No Print / No Varnish  
Lot no. & Exp. date

3 11673 45307 0



**Drug Facts (continued)**

alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions?** Call 1-800-310-6374

**Drug Facts (continued)**

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, croscollon, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, malodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2021 Target Brands, Inc.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING

**PARENTS:**  
Learn what you need to know about  
www.Target.com/parents

Compare to active ingredients in **Maximum Strength Mucinex® SINUS-MAX® Pressure, Pain & Cough\***

**maximum strength  
sinus pressure,  
pain + cough**

**acetaminophen** (pain reliever)  
**dextromethorphan HBr** (cough suppressant)  
**guaifenesin** (expectorant)  
**phenylephrine HCl** (nasal decongestant)

relieves headache,  
sinus pressure and congestion  
controls cough  
thins and loosens mucus

**up&up**

**20 CAPLETS**

**Drug Facts (continued)**

**Uses** ■ temporarily relieves:  
■ sinus congestion and pressure  
■ minor aches and pains  
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the  
bronchial passageways of bothersome mucus and make coughs more  
productive ■ temporarily promotes nasal and/or sinus drainage

**Drug Facts (continued)**

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ blisters ■ rash ■ skin reddening  
If a skin reaction occurs, stop use and seek medical help right away.  
Do not use:  
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ If you have ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)**

**Directions**  
■ do not take more than directed  
■ do not take more than 12 caplets in any 24-hour period  
■ adults and children 12 years and over: take 2 caplets every 4 hours  
■ children under 12 years: do not use

**Other information**  
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR  
BLISTER IS TORN OR BROKEN  
■ see end flap for expiration date and lot number

This product is not manufactured or distributed by R8 Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® SINUS-MAX® Pressure, Pain & Cough. 5/2024 REV02/0461709

No print/No varnish  
Lot & Exp date

094 14 0128 R01 C-001472-01-109




3 11673 71609 0

B-1920-617-09-S  
REV0210A61709

[illegible]







[illegible]

DO NOT USE IF PRINTED NECKBAND  
IS BROKEN OR MISSING

Compare to active ingredients in  
Vicks® DayQuil® Cold & Flu\*

## cold/flu relief multi-symptom day/non-drowsy

pain reliever/fever reducer, cough  
suppressant, nasal decongestant

**acetaminophen** – aches/fever  
dextromethorphan HBr – cough  
phenylephrine HCl – nasal congestion  
alcohol free, antihistamine free



TWO – 12 FL OZ (355 mL), TOTAL – 24 FL OZ (1.5 pt)(710 mL)

NDC 11673-733-02

Compare to active ingredients in  
Vicks® NyQuil® Cold & Flu\*

## cold/flu relief multi-symptom night

antihistamine, cough suppressant,  
fever reducer/pain reliever

**acetaminophen** – aches/fever/sore throat  
dextromethorphan HBr – cough  
doxylamine succinate –  
sneezing/runny nose  
does not contain pseudoephedrine  
**ALCOHOL 10%**



**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Distributed by Target Corporation  
Minneapolis, MN 55403  
©2014 Target Brands, Inc.  
Shop Target.com

\*These products are not manufactured or distributed by Procter & Gamble,  
distributor of Vicks® DayQuil® Cold & Flu and Vicks® NyQuil® Cold & Flu.



Nighttime Cold & Flu							
<b>Drug Facts (continued)</b>  <b>Other information</b> ■ each 30 mL dose cup contains: sodium 39 mg ■ store at 20-25°C (68-77°F)  <b>Inactive ingredients</b> alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate  <b>Questions? Call 1-800-910-6874</b>	<b>Drug Facts</b>  <b>Active ingredients (in each 30 mL dose cup)</b> Acetaminophen 650 mg.....Pain reliever/fever reducer Dextromethorphan HBr 30 mg.....Cough suppressant Doxylamine succinate 12.5 mg.....Antihistamine  <b>Uses</b> temporarily relieves common cold/flu symptoms: ■ headache ■ cough due to minor throat and bronchial irritation ■ sore throat ■ minor aches and pains ■ runny nose and sneezing ■ fever						
<b>Drug Facts (continued)</b>  <b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. The maximum daily dose of this product is 4 doses (2,600 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Sore throat warning:</b> If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  <b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleepy ■ if you have ever had an allergic reaction to this product or any of its ingredients  <b>Ask a doctor before use if you have</b> ■ liver disease ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ a sodium-restricted diet  <b>Ask a doctor or pharmacist before use if you are</b> ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin  <b>When using this product</b> ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness  <b>Stop use and ask a doctor if</b> ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  <b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  <b>Directions</b> ■ take only as directed - see Liver warning ■ use dose cup ■ do not exceed 4 doses per 24 hrs <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL (2 TBSP) every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table> ■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing		adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs						
children 4 to under 12 yrs	ask a doctor						
children under 4 yrs	do not use						



**Drug Facts** (continued)**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 4 doses (adult: 2,600 mg acetaminophen; child: 1,300 mg acetaminophen) in 24 hours. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease ■ high blood pressure ■ heart disease ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

**Stop use and ask a doctor if** ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed - see Liver warning

- use dose cup ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL (2 TBSP) every 4 hrs
children 6 to under 12 yrs	15 mL (1 TBSP) every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

**Drug Facts****Daytime Cold & Flu****Active ingredients (in each 15 mL tablespoon)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Purpose**

- Uses** temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
  - headache
  - minor aches and pains
  - fever
  - sore throat
  - nasal congestion

**Drug Facts** (continued)**Other information**



- each tablespoon contains: sodium 7 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients** butylated hydroxyanisole, edetate disodium, FD&C

yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions? Call 1-800-910-6874**









3106

## Nighttime Cold &amp; Flu

## Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphen HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains
- fever ■ runny nose and sneezing

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

## Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

## When using this product

- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

## Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
  - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- |                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 6 to under 12 yrs      | ask a doctor      |
| children under 6 yrs            | do not use        |

## Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

## Daytime Cold &amp; Flu

## Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphen HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

## Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion ■ cough due to minor throat and bronchial irritation
- sore throat ■ headache ■ minor aches and pains ■ fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
  - child takes more than 5 doses in 24 hours
  - taken with other drugs containing acetaminophen
  - adult has 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

## Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

## When using this product do not use more than directed

## Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
  - pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
  - only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
- |                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

## Other information

- each 15 mL contains: sodium 7 mg ■ store at 20-25°C (68-77°F)

**inactive ingredients:** arcorix, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions? Call 1-888-547-7400**

**inactive ingredients:** butylated hydroxyanisole, edimate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions? Call 1-888-547-7400**



see new warnings

maximum  
strength\*\*

## daytime severe cold

### acetaminophen

(pain reliever/fever reducer)  
dextromethorphan HBr  
(cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl  
(nasal decongestant)

relieves headache,  
fever and sore throat

relieves nasal and  
chest congestion

controls cough

thins and  
loosens mucus

up&up



ACTUAL SIZE

AGES

12+

see new warnings

maximum  
strength\*\*

## nighttime cold and flu

### acetaminophen

(pain reliever/fever reducer)  
diphenhydramine HCl  
(antihistamine)  
phenylephrine HCl  
(nasal decongestant)

relieves headache,  
fever and sore throat

relieves nasal  
congestion

relieves sneezing  
and runny nose

up&up






ACTUAL SIZE

AGES

12+







## Daytime Vaporice™ Cold &amp; Flu

**Active ingredients (in each caplet)**

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms: ■ fever ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ headache ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ itchy nose and sneezing ■ helps

bos en chle

- promotes nasal and/or sinus drainage

## Drug Facts (continued)

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.  
Allergic alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
Some throbbing and pain of the throat is severe, persists for more than  
2 days, is accompanied or followed by fever, headache, rash, nausea,  
or vomiting, can call a doctor promptly.

---

Do not use ■ with any other drug containing acetaminophen  
(prescription or non prescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist. **■** If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emetic conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **■** If you are pregnant or also taking other

Ask a doctor about any use if you have: [www.thirdspace.com](http://www.thirdspace.com) • [www.dofab.com](http://www.dofab.com)

ask a doctor before use if you have: ■ thyroid disease ■ diabetes ■ liver disease ■ heart disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed.

Store away from children and pets. For more information, please see the enclosed information.

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache in a later

These could be signs of a serious condition.

## Drug Facts continued

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:** ■ Take  
■ do not exceed 8 can

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

■ store at 20–25°C (68°/77°F)

3 70030 27574 8

**PARENTS:**  
Learn about test medicine and use  
[www.SmMedsAbuse.com](http://www.SmMedsAbuse.com)

New Facts (continued)

**Inactive ingredients:** croscarmellose sodium, crospovidone, D&C yellow #5 aluminum lake, flux, mabactatin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucrose, etc. (flavoring drops).

Question 2 of 11 988-547-7900